The conference of young scientists «Pidstryhach readings -2020» May 26–28, 2020, Lviv

UDC 512.04

MINIMAL GENERATING SET AND STRUCTURE OF A WREATH PRODUCT OF GROUPS AND THE FUNDAMENTAL GROUP OF AN ORBIT OF MORSE FUNCTION

Ruslan Skuratovskii, Aled Williams

Igor Sikorsky Kiev Polytechnic Institute, r.skuratovskii@kpi.ua Cardiff University, Cardiff, UK, williamsae13@cardiff.ac.uk

The quotient group of the restricted and unrestricted wreath product by its commutator is found. The generic sets of commutator of wreath product were investigated.

We generalize the results presented in the book of Meldrum J. [1] about commutator subgroup of wreath products since, as well as considering regular wreath products, we consider those which are not regular (in the sense that the active group $\mathcal A$ does not have to act faithfully). The fundamental group of orbits of a Morse function $f:M\to\mathbb R$ defined upon a Möbius band M with respect to the right action of the group of diffeomorphisms $\mathcal D(M)$ has been investigated.

Denote the set of all the orbits of \mathcal{A} on X by \mathcal{O} , if this set is finite then by \mathcal{O}_f . Recall that the direct product indexed by infinite set consists of all infinite sequences, and the direct sum consists only of sequences with finitely many elements distinct from zero. Denote by $Z(\tilde{\triangle}(\mathcal{B}))$ the subgroup of diagonal subgroup [2] $Fun(X, Z(\mathcal{B}))$ of functions $f: X \to Z(\mathcal{B})$ which are constant on each orbit of action of A on X for unrestricted wreath product, and denote by $Z(\triangle(\mathcal{B}^n))$ the subgroup of diagonal $Fun(X, Z(\mathcal{B}^n))$ of functions with the same property for restricted wreath product, where n is number of non-trivial coordinates in base of wreath product.

Theorem 1. A centre of the group $(A, X) \wr B$ is direct product of normal closure of centre of a diagonal of $Z(\mathcal{B}^n)$ i.e. $(E \times Z(\triangle(\mathcal{B}^n)))$, trivial an element, and intersection of $(K) \times E$ with Z(A). In other words,

$$Z((\mathcal{A},X) \wr \mathcal{B}) = \langle (1; \underbrace{h,h,\ldots,h}_{n}), e, Z(\mathcal{K},X) \wr \mathcal{E} \rangle \simeq (Z(\mathcal{A}) \cap \mathcal{K}) \times Z(\triangle(\mathcal{B}^{n})),$$

where $h \in Z(\mathcal{B}), |X| = n$.

For restricted wreath product with n non-trivial coordinate: $Z((\mathcal{A}, X) \wr \mathcal{B}) = \langle (1; \ldots, h, h, \ldots, h, \ldots), e, Z(\mathcal{K}, X) \wr \mathcal{E} \rangle \simeq (Z(\mathcal{A}) \cap \mathcal{K}) \times Z(\triangle(\mathcal{B}^n)) \simeq \bigoplus_{j \in O_f} (Z(\mathcal{A}) \cap \mathcal{K}) \times Z(\mathcal{B}).$

The conference of young scientists «Pidstryhach readings – 2020» May 26–28, 2020, Lviv

In case of unrestricted wreath product we have: $Z((\mathcal{A}, X) \wr \mathcal{B}) = \langle (1; \ldots, h_{-1}, h_0, h_1, \ldots, h_i, h_{i+1}, \ldots), e, Z(\mathcal{K}, X) \wr \mathcal{E} \rangle \simeq (Z(\mathcal{A}) \cap \mathcal{K}) \times Z(\tilde{\triangle}(\mathcal{B})) = \prod_{i \in O} (Z(\mathcal{A}) \cap \mathcal{K}) \times Z(\mathcal{B}).$

Theorem 2. If $W = (A, X) \wr (B, Y)$, where |X| = n, |Y| = m and active group A acts on X transitively, then

$$d(G') \le (n-1)d(\mathcal{B}) + d(\mathcal{B}') + d(\mathcal{A}').$$

Theorem 3. The quotient group of a restricted wreath products $G = Z \wr_Z Z$ by a commutator subgroup is isomorphic to $\mathbb{Z} \times \mathbb{Z}$. In previous conditions if $G = A \wr_X B$ then, $G/G' = A/A' \times B/B'$. If $G = Z_n \wr Z_m$, where (m, n) = 1, then d(G/G') = 1. If $G = Z \wr Z$ is an unrestricted regular wreath product then $G/G' \simeq Z \times E \simeq Z$.

- Meldrum John DP. Wreath products of groups and semigroups. volume 74, CRC Press, London, 1995.
- 2. Dixon J.D., Mortimer B. Permutation Groups. Springer-Verlag, New York, 1996.
- 3. Skuratovskii R. The Derived Subgroups of Sylow 2-Subgroups of the Alternating Group and Commutator Width of Wreath Product of Groups // Mathematics, Basel, Switzerland. 2020. 8 (4). P. 3–22.
- Skuratovskii R. V. The commutator subgroup of Sylow 2-subgroups of alternating group, commutator width of wreath product // [Source: Arxiv.org], access mode: https://arxiv.org/pdf/1903.08765.pdf.
- 5. Skuratovskii R. Minimal generating sets for wreath products of cyclic groups, groups of automorphisms of ribe graph and fundumental groups of some morse functions orbits // In Algebra, Topology and Analysis, Odessa, Ukraine. 2016. P. 121–123.
- 6. Skuratovskii R. V. Generating Sets and a Structure of the Wreath Product of Groups with Non-Faithful Group Action // Int. J. Anal. Appl. 2020. 18. P.104–116.

МІНІМАЛЬНА СИСТЕМА ТВІРНИХ І СТРУКТУРА ВІНЦЕВОГО ДОБУТКУ ГРУП І ФУНДАМЕНТАЛЬНА ГРУПА ОРБІТ ФУНКЦІЇ МОРСА

Знайдено фактор групу обмеженого і необмеженого вінцевого добутку по його коммутанту. Досліджено системи твірних комутанта вінцевого добутку. В роботі узагальнено дослідження комутанта вінцевого добутку, що викладені у книзі Мелдрума, на випадок не регулярного вінцевого добутку, де дія активної групи не точна. Фундаментальну групу орбіт функції Морса $f: M \to \mathbb{R}$, що визначена на стрічці Мьобіуса M, досліджено.